

>> SYNTHESIZE MOUNTAINS OF KNOWLEDGE <<

Submitted Abstract

ID IMC22-FSAbstr- 358

First Author First Name Last Name	Dorita (1) Rostkier-Edelstein
Submitting Author First Name Last Name	Dorita Rostkier-Edelstein
Correspondence	dorita.rostkier-edelstein@mail.huji.ac.il
Co-Authors >> E-Mails will be not listed	Kunin, Pavel (2); Alpert, Pinhas (2)
Organisations	1: The Hebrew University of Jerusalem, Israel 2: Tel-Aviv University, Israel
Country	Israel
Region	Middle East
Title	High-Resolution Wrf Simulations Of The Mediterranean Sea Breeze Penetration Into The Dead Sea Valley: Sensitivity Tests And Optimal Model Configuration.
Keywords	Wrf, Sensitivity Tests, Configuration, Parameterizations, High Resolution
Type	List Of Focus Session
Focus Session ID	82

Abstract

The Mediterranean Sea breeze (MSB) is very pronounced during the spring and the summer in the Levant. It flows up the Judean hills, then, it drops into the Dead Sea (DS) valley in the early evening hours. Hence, this flow descends about 1200 m from the Judean Mountains to the DS at about -430 m below MSL. We used WRF model to analyze and better understand the complex processes in and around the DS area, e.g., the MSB structure and its propagation into the DS Valley. A pre-requisite for relying on model simulations to analyze atmospheric processes is to verify their skill with respect to observations. WRF was configured with four nested domain with 30, 10, 3.3 and 1.1 km grid spacing. The model sensitivity was checked for: (1) landuse/vegetation databases, (2) PBL schemes, (3) number of vertical levels, (4) atmospheric and soil initial and lateral boundary conditions from different global models (IC/BC), (5) initial conditions for the finest domain, (7) initialization time of finest domain. Model simulations were verified against unique very high-resolution observations first conducted in the DS valley as part of the Virtual Institute DEad SEa Research Venue (DESERVE) project using the KITcube instruments (a ground-based microwave radiometer, two wind lidars and radiosoundings) along with Energy Balance Stations. These provided horizontal and vertical winds, temperature, humidity, pressure, radiation, and visibility data. Evaluation of model simulations against observations was made for seven parameters that were used to analyze the atmospheric dynamics of the MSB penetration into the DS valley: surface specific humidity, temperature and wind; vertically integrated water vapor; time evolution of horizontal and vertical-wind and profile; and time of MSB arrival to the DS valley. Our sensitivity study shows that the surface variables and the temporal evolution of the horizontal and vertical wind were most sensitive to the PBL scheme and IC/BC from global models. Vertically integrated water vapor was most sensitive to the PBL scheme. The time of arrival of the MSB to the DS valley was most sensitive to IC/BC from global models. The best simulation includes landuse/vegetation from MODIS 15-seconds resolution dataset, the Mellor-YamadaJanjic PBL scheme, 40 vertical levels, IC/BC from NCEP/GFS global model for all nested domains, and later initialization time of the of the finest domain with respect to the coarser ones by 24 hours. In our presentation, we will discuss physical and numerical factors responsible for the most favorable outcomes.